arXiv:1209.2073v1 [math.DS] 10 Sep 2012

NECESSARY AND SUFFICIENT CONDITIONS FOR A DYNAMICAL SYSTEM TO BE TOPOLOGICALLY CONJUGATE TO A GIVEN SUBSTITUTION MINIMAL SYSTEM

ETHAN M. COVEN, ANDREW DYKSTRA, MICHAEL KEANE,
AND MICHELLE LEMASURIER

Abstract. We find necessary and sufficient conditions for a dynamical system to be topologically conjugate to any given substitution minimal system, thus extending the results in [CKL] for the Morse and Toeplitz substitutions.

1. Introduction

In [CKL] three of the authors characterized those symbolic minimal systems that are topologically conjugate to the Morse Minimal System (the closure of the shift-orbit of the famous Morse-Thue sequence) and those topologically conjugate to the closely related Toeplitz Minimal System. The Morse result is that a symbolic minimal system (Y, σ) is topologically conjugate to the Morse system if and only if there exist $N \geq 1$ and 2^N-blocks C_0 and C_1 that appear in Y such that every point in Y can be written as a concatenation of C_0's and C_1's in exactly one way and such that the sequences of C's in these concatenations in some sense "mirror" points in the Morse system.

The arguments in [CKL] are valid only for substitutions sharing some of the properties of the Morse or Toeplitz substitutions. We introduce arguments in Theorem 1 to extend the Morse and Toeplitz results to all substitution minimal systems generated by constant length substitutions. In Theorem 2 we show that the "mirroring" property of the Morse theorem in [CKL] holds for the class of primitive, one-to-one substitutions θ of constant length $L \geq 3$ such that for all $s \neq t$, $\theta(s)$ and $\theta(t)$ disagree at some place other than the first or last.

We show in Theorem 3 that if θ is a primitive substitution of constant length that generates an infinite system (X_θ, σ) and if ζ is a primitive,

---

Date: September 4, 2012.

2010 *Mathematics Subject Classification.* 37B10.

Key words and phrases. Substitution, substitution minimal system, topologically conjugate.

one-to-one substitution of constant length that generates an infinite system (X_ζ, σ) topologically conjugate to (X_θ, σ), then the number of letters in ζ is bounded above by the number of 3-blocks that appear in X_θ. We give an example to show that the bound is attained for the Morse system.

2. Background

In this paper a **dynamical system** is a pair (X, T), where X is a compact metric space and $T : X \to X$ is a homeomorphism. The notion of "sameness" for dynamical systems is **topological conjugacy**: (X, T) and (Y, S) are topologically conjugate iff there exists a homeomorphism $F : (X, T) \to (Y, S)$ such that $F \circ T \equiv S \circ F$. F is called a topological conjugacy and $F^{-1} : (Y, S) \to (X, T)$ is also a topological conjugacy. A **topological semi-conjugacy**, also called a **factor map**, is a continuous, onto map $G : (X, T) \to (Y, S)$ such that $F \circ T \equiv S \circ F$.

A dynamical system (X, T) is called **minimal** iff it contains no proper subsystem. Equivalently, if $X' \subseteq X$ is nonempty, closed, and T-invariant (i.e. $T(X') \subseteq X'$), then $X' = X$. Equivalently again, every orbit $\{T^n(x) : -\infty < n < \infty\}$ is dense in X.

A **symbolic system** is a subsystem of some $(\prod_{-\infty}^{\infty} \mathcal{S}, \sigma)$, where $\mathcal{S}$ is a finite set of symbols and $\sigma : \prod_{-\infty}^{\infty} \mathcal{S} \to \prod_{-\infty}^{\infty} \mathcal{S}$ is the (left) **shift**: $(\sigma(x))_i = x_{i+1}, -\infty < i < \infty$. As is customary, we shall abuse notation and write (X, σ) instead of $(X, \sigma|_X)$. A basic fact about symbolic systems is the Curtis-Hedlund-Lyndon Theorem [LM]: any topological conjugacy or semi-conjugacy $F : (X, \sigma) \to (Y, \sigma)$ between symbolic systems is given by a **local rule** $f : \mathcal{S}_X^{m+1+a} \to \mathcal{S}_Y$, where for every $x \in X$, $(F(x))_i = f(x_{i-m}, \ldots, x_{i+a})$, $-\infty < i < \infty$. Here $m \geq 0$ is called the **memory** and $a \geq 0$ the **anticipation** of F. In this case F is called an $(m + 1 + a)$-block map. Note that by adding superfluous variables an r-block map is also an s-block map for all $s \geq r$.

A **substitution** of constant length $L \geq 2$ is a mapping $\theta : \mathcal{S} \to \mathcal{S}^L$ from a finite set of symbols $\mathcal{S}$ to the set of L-blocks, i.e. blocks of length L, with entries from $\mathcal{S}$. The classic example is the **Morse** substitution: $0 \mapsto 01, 1 \mapsto 10$. A substitution θ can be extended to a mapping of finite blocks by concatenation. For example, $\theta(st) := \theta(s)\theta(t)$. The powers of θ are defined in the obvious way. For example, if $\theta(s) = tuv$, then $\theta^2(s) := \theta(t)\theta(u)\theta(v)$. The symbolic system **generated** by θ is (X_θ, σ), where X_θ is the smallest, closed, σ-invariant set such that for every $s \in \mathcal{S}$ and every $n \geq 1$, $\theta^n(s)$ appears in every point in X_θ. It follows from the definition of X_θ that

Lemma. $X_\theta = X_{\theta^2} = X_{\theta^3} = \dots$

The following lemma shows that when studying symbolic minimal systems up to topological conjugacy, there is no harm in assuming that θ is one-to-one, i.e. if $s \neq t$, then $\theta(s) \neq \theta(t)$.

Lemma. *For any substitution θ of constant length L, there is a one-to-one substitution θ' of constant length L such that (X_θ, σ) is topologically conjugate to $(X_{\theta'}, \sigma)$.*

Proof. If $\theta(s) = \theta(t)$, identify s and t. Continue. □

The following lemma shows that when studying symbolic minimal systems, there is no harm in assuming that θ is **primitive**, i.e. there exists $n \geq 1$ such that for every $s, t \in \mathcal{S}$, s appears in $\theta^n(t)$.

Lemma. [D] *If θ is primitive, then it generates a unique symbolic system and that system is minimal.*

An important property of substitution minimal systems generated by primitive substitutions is **recognizability**, also called **unique decipherability**, defined in the following lemma.

Lemma. [M, Théorèmes 1 and 2] *Let θ be a primitive substitution of constant length that generates an infinite minimal system (X_θ, σ). Then every point in X_θ can be written as a concatenation of the blocks $\theta(s)$ in exactly one way.*

3. Conjugacy to a given substitution minimal system

In this section we find in Theorem 1 necessary and sufficient conditions for a symbolic minimal system to be topologically conjugate to a given substitution minimal system. For a subclass of substitutions, including the Morse substitution, we find in Theorem 2 a result with a simpler statement and a simpler proof than for Theorem 1.

Recall from the Background section that the property of primitivity is natural for substitutions and the property of one-to-oneness is harmless. It turns out that the reasonable property that the substitution minimal system is infinite is needed to make the proofs work.

Theorem 1. *Let θ be a primitive, one-to-one substitution of constant length $L \geq 2$ such that X_θ is infinite, and let (Y, σ) be a symbolic minimal system. Then (Y, σ) is topologically conjugate to (X_θ, σ) if and only if*

(1) *there exist* $N \geq 1$ *and a collection* $\mathcal{B}$ *of* L^N*-blocks such that every point in* Y *can be written as a concatenation of* $\mathcal{B}$*-blocks in exactly one way,*
(2) *there exists a 2-block semiconjugacy* $G : (X_\theta, \sigma) \to (Y_0, \sigma^{L^N})$*, where* Y_0 *is the set of points in* Y *such that the* $\mathcal{B}$*-blocks start at multiples of* L^N*, and*
(3) *if* stu *and* $s't'u'$ *are 3-blocks appearing in* X_θ *and* $t \neq t'$*, then* $g(stu) \neq g(s't'u')$*, where* g *is the local rule of* G*.*

Proof. (ONLY IF) Let $F : (X_\theta, \sigma) \to (Y, \sigma)$ be a topological conjugacy. By composing with a power of the shift, we may assume that F has no memory. Let f be a local rule of F. Choose N so that L^N is greater than the anticipation of F and

$(*)$ for all 3-blocks stu and $s't'u'$ appearing in X_θ with $t \neq t'$, $f(\theta^N(s)\theta^N(t)\theta^N(u)) \neq f(\theta^N(s')\theta^N(t')\theta^N(u'))$.

To see that N can be chosen so that $(*)$ holds, suppose not. Then equality holds for some stu and $s't'u'$ with $t \neq t'$ and infinitely many n. For every n and every s, t, u, $\theta^n(s)\theta^n(t)\theta^n(u)$ can be extended to a doubly infinite sequence, that we also call $\theta^n(s)\theta^n(t)\theta^n(u)$, with the zeroth coordinate coming in a place of disagreement between $\theta^n(t)$ and $\theta^n(t')$. Using the fact that θ and hence every θ^n is one-to-one, there is a subsequence $(\tilde{n})$ of (n) such that both $(\theta^{\tilde{n}}(s)\theta^{\tilde{n}}(t)\theta^{\tilde{n}}(u))$ and $(\theta^{\tilde{n}}(s')\theta^{\tilde{n}}(t')\theta^{\tilde{n}}(u'))$ converge, say to $x \neq x'$. Then $F(x) = F(x')$, contradicting F being one-to-one.

By adding superfluous variables if necessary, we may assume that the anticipation of F is exactly L^N. Define

$$\mathcal{B} := \{f(\theta^N(st)) : st \text{ is a 2-block appearing in } X_\theta\}.$$

We show that condition (1) holds. Since θ and hence θ^N are primitive, every point in $X_{\theta^N} = X_\theta$ can be written as a concatenation of blocks $\theta^N(s)$ in exactly one way [M]. Then, since F is a topological conjugacy, every point in Y can be written as a concatenation of $\mathcal{B}$-blocks in exactly one way.

To show that condition (2) holds, let $G : (X_\theta, \sigma) \to (Y_0, \sigma^{L^N})$, where Y_0 is the set of points in Y such that the $\mathcal{B}$-blocks start at multiples of L^N, be the 2-block semiconjugacy with local rule $g := f \circ \theta^N$.

Condition (3) follows from $(*)$.

(IF) It follows from (3) that the topological semiconjugacy G is one-to-one and hence a topological conjugacy.

Let $X_{\theta,0}$ be the set of points in X_θ such that the blocks $\theta^N(s)$ start at multiples of L^N. Then (X_θ, σ) is topologically conjugate to $(X_{\theta,0}, \sigma^{L^N})$ via the map $s \mapsto \theta^N(s)$.

Standard arguments (e.g. in the proof of the Morse Dynamical Characterization Theorem in [CKL]) show that (X_θ, σ) is topologically conjugate to (Y, σ). □

Now we consider a special case, substitutions θ of length $L \geq 3$ (see Remark below) for which for all $s \neq t$, $\theta(s)$ and $\theta(t)$ disagree in some entry other than the first or last. This class contains the (square of the) Morse substitution.

For such substitutions, condition $(*)$ in the proof of Theorem 1 can be replaced by the stronger condition $(**)$ below and we have the following improvement of Theorem 1 and generalization of the Morse Dynamical Characterization Theorem [CKL].

Theorem 2. *Let θ be a primitive, one-to-one substitution of constant length $L \geq 3$ such that X_θ is infinite, and let (Y, σ) be a symbolic minimal system. Suppose also that any two substitution blocks $\theta(s)$ and $\theta(t)$ with $s \neq t$ disagree somewhere other than in the first or last entry.*

Then (Y, σ) is topologically conjugate to (X_θ, σ) if and only if there exist $N \geq 1$, $a \geq 0$, a collection $\mathcal{B}$ of $(L^N - a)$-blocks that are in one-to-one correspondence with the symbols in X_θ, and a collection $\mathcal{B}'$ of a-blocks such that

1. *every point in Y can be written as a concatenation of alternating $\mathcal{B}$-blocks and $\mathcal{B}'$-blocks in exactly one way,*
2. *the $\mathcal{B}'$-blocks are determined by their nearest $\mathcal{B}$-neighbors,*
3. *for every point in Y, the concatenation of $\mathcal{B}$-blocks "mirrors" a point in X_θ via the one-to-one correspondence above.*

Remark. *We require $L \geq 3$ because condition (3) is vaccuous for $L = 2$. However, this requirement is harmless, for in this case look at θ^2 rather than θ.*

The proof of the ONLY IF direction proceeds as in the proof of Theorem 1, with $F : (X, \sigma) \to (Y, \sigma)$ being a topological conjugacy with no memory and anticipation a and condition $(**)$ below taking the place of condition $(*)$.

$(**)$ there exists N such that L^N is greater than the anticipation of F and such that for all symbols $t \neq t'$ appearing in X_θ, $f(\theta^N(t)) \neq f(\theta^N(t'))$.

Let $\mathcal{B} := \{f(\theta^N(s))\}$ and let $\mathcal{B}'$ be the a-blocks appearing between consecutive $\mathcal{B}$-blocks in points of Y.

To see that condition (2) holds, note that every $\mathcal{B}'$-block appears in some $f(\theta^N(st))$.

The proof of the IF direction is much the same as it is in the proof of Theorem 1.

4. Number of symbols

In this section we prove a relation between the number of symbols in constant length substitutions that generate topologically conjugate systems.

Theorem 3. *Let θ be a primitive, constant length substitution such that X_θ is infinite. Then the number of symbols in a primitive, one-to-one constant length substitution that generates a substitution minimal system topologically conjugate to (X_θ, σ) is at most the number of 3-blocks appearing in X_θ.*

Proof. By [CDL] we may assume that the lengths of θ and ζ are the same. Let $F : (X_\theta, \sigma) \to (X_\zeta, \sigma)$ be a topological conjugacy with no memory and local rule f. Then, with notation as in the proof of Theorem 1, for every symbol s' appearing in X_ζ, $\zeta^N(s')$ is a subblock of some B_1B_2, where $B_1, B_2 \in \mathcal{B}$. But $B_1B_2 = f(\theta^N(stu))$ for some 3-block stu appearing in X_θ. Then we have

$$\#s' = \#\zeta^N(s') \leq \#f(\theta^N(stu)) \leq \#stu,$$

the equality because ζ is one-to-one, and the first inequality because θ and ζ are uniquely decipherable.

□

The following six-symbol example shows that the bound is sharp for the Morse substitution. It is one of the “3-block presentations” of the Morse substitution and so generates a system topologically conjugate to the Morse system (see [CDK]).

$$\zeta(001) = (101)(011)$$
$$\zeta(010) = (110)(100)$$
$$\zeta(011) = (110)(101)$$
$$\zeta(100) = (001)(010)$$
$$\zeta(101) = (001)(011)$$
$$\zeta(110) = (010)(100)$$

With θ the Morse substitution $0 \mapsto 01$, $1 \mapsto 10$, $\zeta(stu) := (s_2t_1t_2)(t_1t_2u_1)$, where $\theta(s) = s_1s_2$, etc.

References

[CDK] Coven, Ethan M., Dekking, F. Michel, and Keane, Michael S., *Substitution minimal sets: conjugacy via N-block substitutions*, in preparation.

[CDL] Coven, Ethan M., Dykstra, Andrew, and LeMasurier, Michelle, *A short proof of a theorem of Cobham on substitutions*, Rocky Mountain J. Math., to appear.

[CKL] Coven, E. M., Keane, M., and LeMasurier, M., *A characterization of the Morse minimal set up to topological conjugacy*, Ergodic Theory Dynamical Systems **28** (2008), 1443–1451.

[D] Dekking, F. M. *Mixing properties of dynamical systems arising from substitutions of constant length*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **42** (1978), 221–239.

[LM] Lind, D. and Marcus, B., *An introduction to symbolic dynamics cand coding*, Cambridge University Press, Cambridge, 1995.

[M] Mossé, B., *Reconnaisabilité des substitutions et complexité des suites automatiques*, Bull. Soc. Math. France **124** (1996), 329–346.

Department of Mathematics, Wesleyan University, Middletown CT 06459
E-mail address: `ecoven@wesleyan.edu`

Department of Mathematics, Hamilton College, Clinton NY 13323
E-mail address: `adykstra@hamilton.edu`

Department of Mathematics, Wesleyan University, Middletown CT 06459
E-mail address: `mkeane@wesleyan.edu`

Department of Mathematics, Hamilton College, Clinton NY 13323
E-mail address: `mlemasur@hamilton.edu`